

Generation mixing phenomena for leptons in e^+e^- collisions

Noriyuki Oshimo

*Department of Physics and Chemistry,
Graduate School of Humanities and Sciences,
Ochanomizu University*

Mixing of quarks or leptons belonging to different generations could give a clue to underlying theories for the generations. We should get phenomenological information about the mixing as much as possible. Until recently most experimental data for the mixing were on the quarks. At present, data on the leptons are accumulating from the neutrino oscillations. However, experiments from different aspects are indispensable for having a deep insight into the lepton generation mixing.

I report the work [NO] which studies measurability of generation mixing for the leptons in e^+e^- collisions within the framework of the supersymmetric standard model (SSM). The SSM is considered a plausible candidate for physics beyond the standard model. This model includes superpartners of quarks and leptons, which could also be mixed among different generations. The interaction of charged leptons, sneutrinos, and charginos can create two different charged leptons with two charginos in e^+e^- annihilation. The same interaction, on the other hand, induces radiative charged-lepton decays, which are constrained by non-observation in experiments to date. Under these constraints, concentrating on the production of e and μ with two lighter charginos, we discuss the possibility of measuring the interaction at e^+e^- colliders.

The mass eigenstates of the leptons or the sneutrinos are generally not the same as their interaction eigenstates. Assuming that no superfield for a right-handed neutrino exists at the electroweak energy scale, the sneutrinos have a 3×3 hermitian mass-squared matrix $\tilde{M}_\nu^2$. The generation mixing of the sneutrinos is traced back to soft supersymmetry-breaking terms. The mass ma-

trix M_l for the charged leptons is proportional to the coefficient matrix of the Higgs couplings. For the left-handed neutrinos, we assume a Majorana mass matrix M_ν . These mass matrices for the leptons are 3×3 . The mass eigenstates are obtained by diagonalizing these matrices as

$$\tilde{U}_\nu^\dagger \tilde{M}_\nu^2 \tilde{U}_\nu = \text{diag}(M_{\tilde{\nu}_1}^2, M_{\tilde{\nu}_2}^2, M_{\tilde{\nu}_3}^2), \quad (1)$$

$$U_{lR}^\dagger M_l U_{lL} = \text{diag}(m_{l_1}, m_{l_2}, m_{l_3}), \quad (2)$$

$$U_\nu^T M_\nu U_\nu = \text{diag}(m_{\nu_1}, m_{\nu_2}, m_{\nu_3}), \quad (3)$$

where $\tilde{U}_\nu$, U_{lR} , U_{lL} , and U_ν are unitary matrices. The masses of the charged leptons e , μ , and τ are denoted by m_{l_1} , m_{l_2} , and m_{l_3} , respectively.

The interaction Lagrangian for charged leptons l_α , sneutrinos $\tilde{\nu}_a$, and charginos ω_i is given by

$$\begin{aligned} \mathcal{L} = & i \frac{g}{\sqrt{2}} (V_C)_{a\alpha} \tilde{\nu}_a^\dagger \overline{\omega_i} \left[\sqrt{2} C_{R1i}^* \left(\frac{1 - \gamma_5}{2} \right) \right. \\ & + C_{L2i}^* \frac{m_{l_\alpha}}{\cos \beta M_W} \left(\frac{1 + \gamma_5}{2} \right) \left. \right] l_\alpha \\ & + \text{H.c.}, \end{aligned} \quad (4)$$

with $V_C = \tilde{U}_\nu^\dagger U_{lL}$. Here, C_L and C_R denote the unitary matrices for diagonalizing the chargino mass matrix which is determined by the SU(2) gaugino mass $\tilde{m}_2$, the Higgsino mass m_H , and the ratio $\tan \beta$ of the vacuum expectation values for the Higgs bosons. The indices α and a for the charged lepton and sneutrino, respectively, represent the generations. The generation mixing is described by V_C , which is a 3×3 unitary matrix. For the independent physical parameters of V_C three mixing angles θ_{12} , θ_{23} , and θ_{13} and one complex phase δ can be taken, the other complex phases being left out by redefinition of particle fields. We adopt the parametrization in the standard form for the Cabibbo-Kobayashi-Maskawa matrix.

A pair of sneutrinos are created in e^+e^- annihilation by the Z -boson and chargino exchange diagrams: $e^+e^- \rightarrow \tilde{\nu}_a\tilde{\nu}_b^*$. Assuming that the sneutrinos are heavier than some of the charginos, the sneutrino can decay into a charged lepton and a chargino $\tilde{\nu}_a^{(*)} \rightarrow l_\alpha^{-(+)}\omega_i^{+(-)}$. The generation mixing could be measured by tagging the charged leptons. However, it is suggested by both theoretical considerations and experimental constraints derived from radiative charged-lepton decays that some sneutrino masses are highly degenerated. Then, the generation mixing has to be analyzed without specifying the sneutrino generations. Our analyses are made under an assumption that the sneutrinos of three generations are produced at the same collision energy and all the sneutrinos can yield a charged lepton of any generation.

We discuss the process $e^+e^- \rightarrow \sum_{a,b} \tilde{\nu}_a\tilde{\nu}_b^* \rightarrow e^-\mu^+\omega_1^+\omega_1^-$. The intermediate sneutrinos are on mass-shell, belonging to any generations. This process shows distinctive final states. The primary leptons e and μ are produced by two-body decays. Their energies are large, unless the mass of the lighter chargino is close to the sneutrino masses. In addition, they have approximately the same value and are monochromatic in each rest frame of the decaying sneutrino. The chargino yields, by three-body decays, two quarks or two leptons with a neutralino. Therefore, the final states involve, in each hemisphere, one charged lepton with a large energy of flat distribution and two jets or one charged lepton, together with large missing energy-momentum. The detection of the process will not be difficult.

The cross section depends on mass differences of the sneutrinos, as the radiative charged-lepton decays do so. As the mass difference between $\tilde{\nu}_1$ and $\tilde{\nu}_2$ becomes large, the cross section increases. The decay width of $\mu \rightarrow e\gamma$ also increases and becomes too large, thus discarding some ranges of the mass difference. The mass of $\tilde{\nu}_3$ does not sensitively affect the cross section nor the decay width. The constraints from the decays $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$ are not very stringent, so that large mass differences between $\tilde{\nu}_1$ and $\tilde{\nu}_3$ and between $\tilde{\nu}_2$ and $\tilde{\nu}_3$ are allowed. The parameters $\tilde{m}_2$ and m_H do not affect much the mass difference between $\tilde{\nu}_1$ and $\tilde{\nu}_2$ allowed by the radiative charged-lepton

decays. However, the cross section varies manifestly with these parameters, depending on the relative magnitudes of $\tilde{m}_2$ and m_H . For a smaller magnitude of $\tilde{m}_2/m_H$, the SU(2)-gaugino component of the lighter chargino is larger. Then, the cross section of $e^+e^- \rightarrow \sum_{a,b} \tilde{\nu}_a\tilde{\nu}_b^*$ and thus that of $e^+e^- \rightarrow e^-\mu^+\omega_1^+\omega_1^-$ increase.

The dependencies of the cross section on the mixing angles of V_C and $\tan\beta$ are also manifest. As the mixing angle increases, the allowed ranges for the mass difference between $\tilde{\nu}_1$ and $\tilde{\nu}_2$ become narrow, while the cross section becomes large. These angle dependencies are primarily determined by θ_{12} . Different values for θ_{23} and θ_{13} do not alter much the cross section and the allowed mass difference, as long as these mixing angles are small. For a larger value of $\tan\beta$, the allowed region becomes small, though the cross section does not vary much with it.

Under the constraints from the presently available experiments, the cross section of $e^+e^- \rightarrow e^-\mu^+\omega_1^+\omega_1^-$ is larger than 0.05 fb in sizable regions of the parameter space. It should be noted that there also exists a charge conjugate process which has the same cross section. For the integrated luminosity of 100 fb $^{-1}$, more than ten events are expected there. Although possible backgrounds have to be taken into account for estimating realistically available events, a number of order of ten would not be insufficient for detection in near future experiments.

In summary, the examination of the process $e^+e^- \rightarrow e^-\mu^+\omega_1^+\omega_1^-$ will enable us to explore the generation mixing peculiar to the SSM. Moreover, the SSM parameters for the chargino and neutralino sector affect non-trivially its cross section. The study of the generation mixing will give information on these SSM parameters.

This work is supported in part by the Grant-in-Aid for Scientific Research on Priority Areas (No. 16028204) from the Ministry of Education, Science and Culture, Japan.

References

[NO] N. Oshimo, Euro. Phys. J **C 39** (2005), 383.